

ON THE STRONG CHROMATIC INDEX AND MAXIMUM INDUCED MATCHING OF TREE-COGRAPHS, PERMUTATION GRAPHS AND CHORDAL BIPARTITE GRAPHS

Ton Kloks^{*1}, Chin-Ting Ung¹, and Yue-Li Wang²

¹ Department of Computer Science
National Tsing Hua University, No. 101, Sec. 2, Kuang Fu Rd., Hsinchu, Taiwan
wonderboy0915@gmail.com

² Department of Information Management
National Taiwan University of Science and Technology
No. 43, Sec. 4, Keelung Rd., Taipei, 106, Taiwan
ylwang@cs.ntust.edu.tw

Abstract. We show that there exist linear-time algorithms that compute the strong chromatic index and a maximum induced matching of tree-cographs when the decomposition tree is a part of the input. We also show that there exist efficient algorithms for the strong chromatic index of (bi-partite) permutation graphs and of chordal bipartite graphs.

1 Introduction

Definition 1. Let $G = (V, E)$ be a graph. A strong edge coloring of G is a proper edge coloring such that no edge is adjacent to two edges of the same color.

Equivalently, a strong edge coloring of G is a vertex coloring of $L(G)^2$, the square of the linegraph of G . The strong chromatic index of G is the minimal integer k such that G has a strong edge coloring with k colors. We denote the strong chromatic index of G by $s\chi'(G)$.

The class of tree-cographs was introduced by Tinhofer in [33].

Definition 2. Tree-cographs are defined recursively by the following rules.

1. Every tree is a tree-cograph.
2. If G is a tree-cograph then also the complement $\bar{G}$ of G is a tree-cograph.
3. For $k \geq 2$, if $G_1, \dots, G_k$ are connected tree-cographs then also the disjoint union is a tree-cograph.

Let G be a tree-cograph. A decomposition tree for G consists of a rooted binary tree T in which each internal node, including the root, is labeled as a join node $\otimes$ or a union node $\oplus$. The leaves of T are labeled by trees or complements of trees. It is easy to see that a decomposition tree for a tree-cograph G can be obtained in $O(n^3)$ time.

* National Science Council of Taiwan Support Grant NSC 99-2218-E-007-016.

2 The strong chromatic index of tree-cographs

The linegraph $L(G)$ of a graph G is the intersection graph of the edges of G [4]. It is well-known that, when G is a tree then the linegraph $L(G)$ of G is a claw-free blockgraph [18]. A graph is chordal if it has no induced cycles of length more than three [12]. Notice that blockgraphs are chordal.

A vertex x in a graph G is simplicial if its neighborhood $N(x)$ induces a clique in G . Chordal graphs are characterized by the property of having a perfect elimination ordering, which is an ordering $[v_1, \dots, v_n]$ of the vertices of G such that v_i is simplicial in the graph induced by $\{v_i, \dots, v_n\}$. A perfect elimination ordering of a chordal graph can be computed in linear time [31]. This implies that chordal graphs have at most n maximal cliques, and the clique number can be computed in linear time.

Theorem 1 ([6]). *If G is a chordal graph then $L(G)^2$ is also chordal.*

Proof. Any chordal graph is the intersection graph of a collection of subtrees of a tree. Let G be the intersection graph of a collection of subtrees of a tree. An intersection model for $L(G)^2$ is obtained by taking the union of every pair of intersecting subtrees. $\square$

Theorem 2 ([8]). *Let $k \in \mathbb{N}$ and let $k \geq 4$. Let G be a graph and assume that G has no induced cycles of length at least k . Then $L(G)^2$ has no induced cycles of length at least k .*

Lemma 1. *Tree-cographs have no induced cycles of length more than four.*

Proof. Let G be a tree-cograph.

First observe that trees are bipartite. It follows that complements of trees have no induced cycles of length more than four.

We prove the claim by induction on the depth of a decomposition tree for G . If G is the union of two tree-cographs G_1 and G_2 then the claim follows by induction since any induced cycle is contained in one of G_1 and G_2 . Assume G is the join of two tree-cographs G_1 and G_2 . Assume that G has an induced cycle C of length at least five. We may assume that C has at least one vertex in each of G_1 and G_2 . If one of G_1 and G_2 has more than two vertices of C , then C has a vertex of degree at least three, which is a contradiction. $\square$

Lemma 2. *Let T be a tree. Then $L(\bar{T})^2$ is a clique.*

Proof. Consider two non-edges $\{a, b\}$ and $\{p, q\}$ of T . If the non-edges share an endpoint then they are adjacent in $L(\bar{T})^2$ since they are already adjacent in $L(\bar{T})$. Otherwise, since T is a tree, at least one pair of $\{a, p\}$, $\{a, q\}$, $\{b, p\}$ and $\{b, q\}$ is a non-edge in T , otherwise T has a 4-cycle. By definition, $\{a, b\}$ and $\{p, q\}$ are adjacent in $L(\bar{T})^2$. $\square$

If G is the union of two tree-cographs G_1 and G_2 then the maximal cardinality of a clique in $L(G)^2$ is, simply, the maximum over the clique numbers of $L(G_1)^2$ and $L(G_2)^2$. The following lemma deals with the join of two tree-cographs.

Lemma 3. Let P and Q be tree-cographs and let G be the join of P and Q . Let X be the set of edges that have one endpoint in P and one endpoint in Q . Then

- (a) X forms a clique in $L(G)^2$,
- (b) every edge of X is adjacent in $L(G)^2$ to every edge of P and to every edge of Q ,
and
- (c) every edge of P is adjacent in $L(G)^2$ to every edge of Q .

Proof. This is an immediate consequence of the definitions. $\square$

For $k \geq 3$, a k -sun is a graph which consists of a clique with k vertices and an independent set with k vertices. There exist orderings $c_1, \dots, c_k$ and $s_1, \dots, s_k$ of the vertices in the clique and independent set such that each s_i is adjacent to c_i and to c_{i+1} for $i = 1, \dots, k-1$ and such that s_k is adjacent to c_k and c_1 . A graph is strongly chordal if it is chordal and has no k -sun, for $k \geq 3$ [13].




Fig. 1. A 3-sun, a gem and a claw

Lemma 4. Let T be a tree. Then $L(T)^2$ is strongly chordal.

Proof. When T is a tree then $L(T)$ is a blockgraph. Obviously, blockgraphs are strongly chordal. Lubiw proves in [29] that all powers of strongly chordal graphs are strongly chordal. $\square$

We strengthen the result of Lemma 4 as follows. Ptolemaic graphs are graphs that are both distance hereditary and chordal [21]. Ptolemaic graphs are gem-free chordal graphs. The following theorem characterizes ptolemaic graphs.

Theorem 3 ([21]). A connected graph is ptolemaic if and only if for all pairs of maximal cliques C_1 and C_2 with $C_1 \cap C_2 \neq \emptyset$, the intersection $C_1 \cap C_2$ separates $C_1 \setminus C_2$ from $C_2 \setminus C_1$.

Lemma 5. Let T be a tree. Then $L(T)^2$ is ptolemaic.

Proof. Consider $L(T)$. Let C be a block and let P and Q be two blocks that each intersect C in one vertex. Since $L(T)$ is claw-free, the intersections of $P \cap C$ and $Q \cap C$ are distinct vertices. The intersection of the maximal cliques $P \cup C$ and $Q \cup C$, which is C , separates $P \setminus Q$ and $Q \setminus P$ in $L(T)^2$. Since all intersecting pairs of maximal cliques are of this form, this proves the lemma. $\square$

Corollary 1. *Let G be a tree-cograph. Then $L(G)^2$ has a decomposition tree with internal nodes labeled as join nodes and union nodes and where the leaves are labeled as ptolemaic graphs.*

From Corollary 1 it follows that $L(G)^2$ is perfect [10], that is, $L(G)^2$ has no odd holes or odd antiholes [28]. This implies that the chromatic number of $L(G)^2$ is equal to the clique number. Therefore, to compute the strong chromatic index of a tree-cograph G it suffices to compute the clique number of $L(G)^2$.

Theorem 4. *Let G be a tree-cograph and let T be a decomposition tree for G . There exists a linear-time algorithm that computes the strong chromatic index of G .*

Proof. First assume that $G = (V, E)$ is a tree. Then the strong chromatic index of G is

$$s\chi'(G) = \max \{ d(x) + d(y) - 1 \mid (x, y) \in E \} \quad (1)$$

where $d(x)$ is the degree of the vertex x . To see this notice that Formula (1) gives the clique number of $L(G)^2$.

Assume that G is the complement of a tree. By Lemma 2 the strong chromatic index is the number of nonedges in G , which is

$$s\chi'(G) = \binom{n}{2} - (n - 1).$$

Assume that G is the union of two tree-cographs G_1 and G_2 . Then, obviously,

$$s\chi'(G) = \max \{ s\chi'(G_1), s\chi'(G_2) \}.$$

Finally, assume that G is the join of two tree-cographs G_1 and G_2 . Let X be the set of edges of G that have one endpoint in G_1 and the other in G_2 . Then, by Lemma 3, we have

$$s\chi'(G) = |X| + s\chi'(G_1) + s\chi'(G_2).$$

The decomposition tree for G has $O(n)$ nodes. For the trees the strong chromatic index can be computed in linear time. In all other cases, the evaluation of $s\chi'(G)$ takes constant time. It follows that this algorithm runs in $O(n)$ time, when a decomposition tree is a part of the input. $\square$

3 Induced matching in tree-cographs

Consider a strong edge coloring of a graph G . Then each color class is an induced matching in G , which is an independent set in $L(G)^2$ [6]. In this section we show that the maximal value of an induced matching in G can be computed in linear time. Again, we assume that a decomposition tree is a part of the input.

Theorem 5. *Let G be a tree-cograph and let T be a decomposition tree for G . Then the maximal number of edges in an induced matching in G can be computed in linear time.*

Proof. In this proof we denote the maximal cardinality of an induced matching in a graph G by $iv(G)$.

First assume that G is a tree. Since the maximum induced matching problem can be formulated in monadic second-order logic, there exists a linear-time algorithm to compute the maximal cardinality of an induced matching in G .

Assume that G is the complement of a tree. By Lemma 2 $L(G)^2$ is a clique. Thus the maximal cardinality of an induced matching in G is one if G has a nonedge and otherwise it is zero.

Assume that G is the union of two tree-cographs G_1 and G_2 . Then

$$iv(G) = iv(G_1) + iv(G_2).$$

Assume that G is the join of two tree-cographs G_1 and G_2 . Then

$$iv(G) = \max \{ iv(G_1), iv(G_2), 1 \}.$$

This proves the theorem. □

4 Permutation graphs

A permutation diagram on n points is obtained as follows. Consider two horizontal lines L_1 and L_2 in the Euclidean plane. For each line L_i consider a linear ordering $\prec_i$ of $\{1, \dots, n\}$ and put points $1, \dots, n$ on L_i in this order. For $k = 1, \dots, n$ connect the two points with the label k by a straight line segment.

Definition 3 ([15]). *A graph G is a permutation graph if it is the intersection graph of the line segments in a permutation diagram.*




Fig. 2. A permutation graph and a permutation diagram

Consider two horizontal lines L_1 and L_2 and on each line L_i choose n intervals. Connect the left - and right endpoint of the k^{th} interval on L_1 with the left

- and right endpoint of the k^{th} interval on L_2 . Thus we obtain a collection of n trapezoids. We call this a trapezoid diagram.

Definition 4. A graph is a trapezoid graph if it is the intersection graph of a collection of trapezoids in a trapezoid diagram.

Lemma 6. If G is a permutation graph then $L(G)^2$ is a trapezoid graph.

Proof. Consider a permutation diagram for G . Each edge of G corresponds to two intersecting line segments in the diagram. The four endpoints of a pair of intersecting line segments define a trapezoid. Two vertices in $L(G)^2$ are adjacent exactly when the corresponding trapezoids intersect (see Proposition 1 in [7]). $\square$

Theorem 6. There exists an $O(n^4)$ algorithm that computes a strong edge coloring in permutation graphs.

Proof. Dagan, et al., [11] show that a trapezoid graph can be colored by a greedy coloring algorithm. It is easy to see that this algorithm can be adapted so that it finds a strong edge-coloring in permutation graphs. $\square$

Remark 1. A somewhat faster coloring algorithm for trapezoid graphs appears in [14]. Their algorithm runs in $O(n \log n)$ time where n is the number of vertices in the trapezoid graph. An adaption of their algorithm yields a strong edge coloring for permutation graphs that runs in $O(m \log n)$ time, where n and m are the number of vertices and edges in the permutation graph.

4.1 Bipartite permutation graphs

Let $G = (A, B, E)$ be a bipartite permutation graph with color classes A and B .

Lemma 7. Let G be a bipartite permutation graph. Then $L(G)^2$ is an interval graph.

Proof. We first show that $L(G)^2$ is chordal. We may assume that $L(G)^2$ is connected.

Let x and y be two non-adjacent vertices in a graph H . An x, y -separator is a set S of vertices which separates x and y in distinct components. The separator is a minimal x, y -separator if no proper subset of S separates x and y . A set S is a minimal separator if there exist non-adjacent vertices x and y such that S is a minimal x, y -separator. Recall that Dirac characterizes chordal graphs by the property that every minimal separator is a clique [12].

Consider the trapezoid diagram. Let S be a minimal separator in the trapezoid graph $L(G)^2$ and consider removing the trapezoids that are in S from the diagram. Every component of $L(G)^2 - S$ is a connected part in the diagram. Consider the left-to-right ordering of the components in the diagram. Since S is a minimal separator there must exist two consecutive components C_1 and C_2 such that every vertex of S has a neighbor in both C_1 and C_2 [5].

Assume that S has two non-adjacent trapezoids t_1 and t_2 . Each of t_i is characterized by two crossing line segments $\{a_i, b_i\}$ of the permutation diagram. Since t_1 and t_2 are not adjacent, any pair of line-segments with one element in $\{a_1, b_1\}$ and the other element in $\{a_2, b_2\}$ are parallel.

Each trapezoid t_i intersects each component C_1 and C_2 . Since pairs of line-segments are parallel, we have that, for some $i \in \{1, 2\}$

- (1) $N_G(a_i) \cap C_1 \subseteq N_G(a_{3-i}) \cap C_1$,
- (2) $N_G(a_i) \cap C_1 \subseteq N_G(b_{3-i}) \cap C_1$,
- (3) $N_G(b_i) \cap C_1 \subseteq N_G(a_{3-i}) \cap C_1$ and
- (4) $N_G(b_i) \cap C_1 \subseteq N_G(b_{3-i}) \cap C_1$,

and the reverse inequalities hold for C_2 . Each trapezoid t_i has at least one line segment of $\{a_i, b_i\}$ intersecting with a line segment of C_i . By the neighborhood containments this implies that G has a triangle, which contradicts that G is bipartite. This proves that S is a clique and by Dirac's characterization $L(G)^2$ is chordal.

Lekkerkerker and Boland prove in [27] that a graph H is an interval graph if and only if H is chordal and H has no asteroidal triple. It is easy to see that a permutation graph has no asteroidal triple [23]. Cameron and Chang prove in [7, 9] that $L(H)^2$ is AT-free whenever a graph H is AT-free. Thus, since $L(G)^2$ is chordal and AT-free, $L(G)^2$ is an interval graph.

This proves the lemma. $\square$

Chang proves in [9] that there exists a linear-time algorithm that computes a maximum induced matching in bipartite permutation graphs. We show that there is a simple linear-time algorithm that computes the strong chromatic index of bipartite permutation graphs.

Theorem 7. *There exists a linear-time algorithm that computes the strong chromatic index of bipartite permutation graphs.*

Proof. Let $G = (A, B, E)$ be a bipartite permutation graph and consider a permutation diagram for G . Let $[a_1, \dots, a_s]$ and $[b_1, \dots, b_t]$ be left-to-right orderings of the vertices of A and B on the topline of the diagram. Assume that a_1 is the left-most endpoint of a line segment on the topline. We may assume that the line segment of a_1 intersects the line segment of b_1 .

Consider the set of edges in the maximal complete bipartite subgraph M in G that consists of the following vertices.

- (a) M contains a_1 and b_1 ,
- (b) M contains all the vertices of A of which the endpoint on the topline is to the left of b_1 ,
- (c) M contains all the vertices of B of which the endpoint on the bottom line is to the left of a_1 .

Notice that M is the set of edges in the complete bipartite subgraph in G induced by

$$N[a_1] \cup N[b_1].$$

Extend the set of edges in M with the edges that have one endpoint in

$$N(a_1) \cup N(b_1).$$

In the diagram this means that we add the line segments that cross one of the line segments corresponding to the vertices of M mentioned above. Call the set of edges in M plus the edges with one endpoint in $N(a_1) \cup N(b_1)$ the extension $\bar{M}$ of M . Notice that $\bar{M}$ is the unique maximal clique that contains the simplicial edge $\{a_1, b_1\}$ in $L(G)^2$.

The second maximal clique in $L(G)^2$ is found by the process described above for the line segments induced by $A \setminus \{a_1\} \cup B$. Likewise, the third maximal clique in $L(G)^2$ is found by repeating the process for the line segments induced by $A \cup B \setminus \{b_1\}$.

Next, remove the vertices a_1 and b_1 and repeat the three steps described above. It is easy to see that the list obtained in this manner contains all the maximal cliques of $L(G)^2$. Notice also that this algorithm can be implemented to run in linear time. Since $L(G)^2$ is perfect the chromatic number is equal to the clique number, so it suffices to keep track of the maximal cardinality of the cliques that are found in the process described above. $\square$

5 Chordal bipartite graphs

Definition 5 ([16, 22]). A bipartite graph is chordal bipartite if it has no induced cycles of length more than four.

A graph is weakly chordal if it has no induced cycle of length more than four or the complement of such a cycle [19]. Weakly chordal graphs are perfect. Notice that chordal bipartite graphs are weakly chordal.

Cameron and Chang prove in [8, 9] that $L(G)^2$ is weakly chordal whenever G is weakly chordal. Thus, if G is chordal bipartite then $L(G)^2$ is perfect and so, in order to compute the strong chromatic index of G it is sufficient to compute the clique number in $L(G)^2$ (see also [1]).

It is well-known that the clique number of a perfect graph can be computed in polynomial time [17].³ The algorithm presented in [20] to compute the clique number of weakly chordal graphs runs in $O(n^3)$ time, where n is the number of vertices in the graph.

³ Actually, this paper shows that for any graph G with $\omega(G) = \chi(G)$ the values of these parameters can be determined in polynomial time. The reason is that Lovász' bound $\vartheta(G)$ for the Shannon capacity of a graph can be computed in polynomial time for all graphs, via the ellipsoid method, and the parameter $\vartheta(G)$ is sandwiched between $\omega(G)$ and $\chi(G)$.

A direct application of their algorithm to solve the strong chromatic index of chordal bipartite graphs G involves computing the graph $L(G)^2$. This graph has m vertices, where m is the number of edges in G . This gives a timebound $O(n^6)$ for computing the strong chromatic index of a chordal bipartite graph (see also [1]).

In this section we show that there is a more efficient method.

Definition 6. A bipartite graph is a chain graph if it has no induced $2K_2$.

Equivalently, a bipartite graph $G = (A, B, E)$ is a chain graph if for every pair x and y in A $N(x) \subseteq N(y)$ or $N(y) \subseteq N(x)$. Notice that a set of edges in G form a clique in $L(G)^2$ exactly when it is a set of edges in an induced chain graph of G . Thus, computing the clique number of $L(G)^2$ is equivalent to finding the maximal number of edges of an induced chain graph of G .

Theorem 8. There exists an $O(n^4)$ algorithm that computes the strong chromatic index of chordal bipartite graphs.

Proof. We use the following property which holds for chordal bipartite graphs $G = (A, B, E)$.

Let x, y and z be three vertices of A . Then at least one of $N(x)$, $N(y)$ and $N(z)$ contains the intersection of the other two [2, Proposition 3].

It follows that the number of nonempty, minimal intersections of neighborhoods of vertices in A is $O(n^2)$. We remark that this can also be seen as follows. First, consider two vertices x and y in A with $N(x) \cap N(y) \neq \emptyset$. Consider adding a vertex z to A with $N(z) = N(x) \cap N(y)$. Then the new graph is again chordal bipartite [3, Lemma 2.3]. Assume that G has no isolated vertex and that no two vertices in A have exactly the same set of neighbors. Then the number of vertices in A is at most $\binom{|B|+1}{2}$ (see, e.g., [25, 26]).

Consider a sequence of intersections $\Omega_1, \dots, \Omega_k$ of neighborhoods of vertices in A , where

$$\Omega_1 \neq \emptyset \quad \text{and} \quad \Omega_1 \subset \dots \subset \Omega_k.$$

That is, each Ω_i is properly contained in Ω_{i+1} , for $i = 1, \dots, k-1$. Observe that we may assume that there exists a vertex x such that $N(x) = \Omega_k$. For $i = 1, \dots, k$ let

$$\Gamma_i = \{x \in A \mid \Omega_i \subseteq N(x)\}.$$

Define,

- (1) $\Omega_1^* = \Omega_1$, and
- (2) for $i > 1$, $\Omega_i^* = \Omega_i \setminus \Omega_{i-1}$.

Then the number of edges in the chain graph defined by $\Omega_1, \dots, \Omega_k$ is

$$\sum_{i=1}^k |\Omega_i^*| \cdot |\Gamma_i|.$$

It is easy to see that the chain with a maximal number of edges can be determined by dynamic programming in $O(n^4)$ time.

This proves the theorem. $\square$

References

1. Abueida, A., A. Busch and R. Sritharan, A min-max property of chordal bipartite graphs with applications, *Graphs and Combinatorics* **26** (2010), pp. 301–313.
2. Acharya, B. and M. Las Vergnas, Hypergraphs with cyclomatic number zero, triangulated graphs, and an inequality, *Journal of Combinatorial Theory, Series B* **33** (1982), pp. 52–56.
3. Anstee, R., Hypergraphs with no special cycles, *Combinatorica* **3** (1983), pp. 141–146.
4. Beineke, L., Characterization of derived graphs, *Journal of Combinatorial theory* **9** (1970), pp. 129–135.
5. Bodlaender, H., T. Kloks, D. Kratsch and H. Müller, Treewidth and minimum fill-in of d-trapezoid graphs, *Journal of Graph Algorithms and Applications* **2** (1998), pp. 1–23.
6. Cameron, K., Induced matchings, *Discrete Applied Mathematics* **24** (1989), pp. 97–102.
7. Cameron, K., Induced matchings in intersection graphs, *Discrete Mathematics* **278** (2004), pp. 1–9.
8. Cameron, K., R. Sritharan, and Y. Tang, Finding a maximum induced matching in weakly chordal graphs, *Discrete Mathematics* **266** (2003), pp. 133–142.
9. Chang, J., Induced matchings in asteroidal triple-free graphs, *Discrete Applied Mathematics* **132** (2004), pp. 67–78.
10. Chudnovsky, M., N. Robertson, P. Seymour and R. Thomas, The strong perfect graph theorem, *Annals of Mathematics* **164** (2006), pp. 51–229.
11. Dagan, I., M. Golumbic and R. Pinter, Trapezoid graphs and their coloring, *Discrete Applied Mathematics* **21** (1988), pp. 35–46.
12. Dirac, G., On rigid circuit graphs, *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg* **25** (1961), pp. 71–76.
13. Farber, M., Characterizations of strongly chordal graphs, *Discrete Mathematics* **43** (1983), pp. 173–189.
14. Felsner, S., R. Müller and L. Wernisch, Trapezoid graphs and generalizations, geometry and algorithms, *Discrete Applied mathematics* **74** (1997), pp. 13–22.
15. Golumbic, M., *Algorithmic graph theory and perfect graphs*, Elsevier, Annals of Discrete Mathematics **57**, 2004.
16. Golumbic, M. and C. Goss, Perfect elimination and chordal bipartite graphs, *Journal of Graph Theory* **2** (1978), pp. 155–163.
17. Grötschel, M., L. Lovász and A. Schrijver, Polynomial algorithms for perfect graphs. In *Annals of Discrete Mathematics*, Elsevier (1984), pp. 325–356.
18. Harary, F., A characterization of block-graphs, *Canadian Mathematical Bulletin* **6** (1963), pp. 1–6.
19. Hayward, R., Weakly triangulated graphs, *Journal of Combinatorial Theory, Series B* **39** (1985), pp. 200–208.
20. Hayward, R., J. Spinrad and R. Sritharan, Improved algorithms for weakly chordal graph, *ACM Transactions on Algorithms* **3** (2007), Article No. 14.
21. Howorka, E., A characterization of ptolemaic graphs, *Journal of Graph Theory* **5** (1981), pp. 323–331.
22. Huang, J., Representation characterizations of chordal bipartite graphs, *Journal of Combinatorial Theory, Series B* **96** (2006), pp. 673–683.
23. Kloks, T., *Treewidth – Computations and approximations*, Springer, Lecture Notes in Computer Science **842**, 1994.

24. Kloks, T. and D. Kratsch, Computing a perfect edge without vertex elimination ordering, *Information Processing Letters* **55** (1995), pp. 11–16.
25. Kloks, T., C. Liu and S. Poon, Feedback vertex set on chordal bipartite graphs. ArXiv: 1104.3915, 2011.
26. Lehel, J., A characterization of totally balanced hypergraphs, *Discrete Mathematics* **57** (1985), pp. 59–65.
27. Lekkerkerker, C. and D. Boland, Representation of finite graphs by a set of intervals on the real line, *Fundamenta Mathematicae* **51** (1962), pp. 45–64.
28. Lovász, L., Perfect graphs. In, L. Beineke and R. Wilson eds., *Selected topics in Graph Theory, Vol. 2*, Academic Press, 1983, pp. 55–87.
29. Lubiw, A., Γ -free matrices. Master’s Thesis, Department of Combinatorics and Optimization, University of Waterloo, Canada, 1982.
30. Lubiw, A., Doubly lexical orderings of matrices, *SIAM Journal on Computing* **16** (1987), pp. 854–879.
31. Rose, D., G. Luecker and R. Tarjan, Algorithmic aspects of vertex elimination on graphs, *SIAM Journal on Computing* **5** (1976), pp. 266–283.
32. Sawada, J. and J. Spinrad, From a simple elimination ordering to a strong elimination ordering in linear time, *Information Processing Letters* **86** (2003), pp. 299–302.
33. Tinhofer, G., Strong tree-cographs are Birkhoff graphs, *Discrete Applied Mathematics* **22** (1989), pp. 275–288.